
AD-A192 755

EXCITED-STATE ENERGETICS AND DYNAMICS OF LARGE
MOLECULES COMPLEXES AND CL.. (U) TEL-AVIV UNIV (ISRAEL)
DEPT OF CHEMISTRY J JORTHOR ET AL. APR 85

UNCLASSIFIED R/D-4759-CH-01 DAJA45-85-C-0008 F/G 7/4 NL



TOTAL PROPERTY METALOGICAL DESCRIPTION OF THE PROPERTY OF THE



OTIC FILE CORY

AD-A192 755

4759- CH-01

FIRST INTERIM REPORT

Contract No. DAJA45-85-C-0008

PROFESSOR JOSHUA JORTNER

School of Chemistry Tel Aviv University Ramat Aviv Campus 69978 Tel Aviv Israel



to character and been approved to paid to release and sales in the paid of the

8 3 23 12 4

PROGRESS REPORT FOR THE PERIOD FEBRUARY - APRIL, 1985

1st Periodic Report

1. Title: EXCITED-STATE ENERGETICS AND DYNAMICS

OF LARGE MOLECULES, COMPLEXES AND

CLUSTERS

2. Principal Investigator: Professor Joshua Jortner

3. Associated Investigator: Professor Uzi Even

4. Contractor: The Department of Chemistry

Tel Aviv University

69978 Tel Aviv

Israel

5. Contract No. DAJA45-85-C-0008

The Research reported in this document has been made possible through the support and sponsorship of the U.S. Government through its European Research Office of the U.S. Army. This report is intended only for the internal management use of the Contractor and the

U.S. Covernment.


STATESTER INTERCACE (IN POSSESSEE IN POSSESSEE IN PROPERTY IN PROPERTY IN PROPERTY IN PROPERTY IN PROPERTY IN

6. PROGRESS IN TECHNICAL APPROACH

Vacuum Ultraviolet Absorption Spectroscopy in Supersonic Expansions. We have combined the techniques of vacuum ultraviolet (VUV) spectroscopy together with planar supersonic jets, which allows for the interrogation of absorption spectra of large molecules cooled in supersonic expansions in the near VUV region. The experimental setup consists of a high pressure Xe lamp, car optics, a vacuum ultraviolet spectrograph and a nozzle slit (0.27x90 mm, repetition rate 9 Hz and gas pulse duration 300 µsec). The characteristics of this spectroscopic setup are: (i) Energy range 6-10 eV. (ii) Spectral resolution 0.14. (iii) Routine measurements of high-energy absorption spectra. (iv) Interrogation of fluorescence excitation spectra of the parent molecule or its photoproducts with limiting quantum yields of Y > 10-4.

7. ACCOMPLISHMENTS OF OBJECTIVES

- 7.A Energetics of Rydberg States of Jet Cooled Molecules. VUV absorption spectra of benzene, benzene-D₆ and naphthalene cooled in planar supersonic expansions were measured over the range 2000-160**0**Å, providing evidence on energetics, line broadening and interference effects.
- 7.B Intramolecular Relaxation of Rydberg States. Information on intramolecular dynamics of extravalence excitations of benzene was obtained from lineshape analysis. The lineshape of the $^{3}P_{XY}(0)$ Rydberg is Lorentzian, whose homogeneous width result in the lifetime $\tau = 0.19 \pm 0.02$ psec for C_6H_6 and $\tau \simeq 0.22 \pm 0.02$ psec for C_6D_6 , implying that (i) the relaxation of the Rydberg is characterized by moderate energy gaps, and (ii) the electronic relaxation rate of the Rydberg is considerably less efficient than that of the intravalence excitation in the same energy domain.





8. PUBLICATIONS

- A. Amirav and J. Jortner Vacuum Ultraviolet Absorption Spectroscopy in Supersonic Expansions J. Chem. Phys. (Communication) (in press)
- (2) U. Even, Z. Berkovitch-Yellin and J. Jortner Electronic Excitations of the Free-Base Porphine-Ar van der Waals Complex Canad. J. Chem. (C. Sandorfy-Special Issue) (in press)

END DATE FILMED 6-1988 DTIC